

POSTER PRESENTATION

Open Access

P02.64. Vitamin D sufficiency is necessary for integrative treatment-associated improvements in chronic pain status

G Plotnikoff*, J Dusek, BraveNet Practice Based Research Network

From International Research Congress on Integrative Medicine and Health 2012
 Portland, Oregon, USA. 15-18 May 2012

Purpose

To examine the relationship between improved vitamin D status and reported severity of chronic pain.

Methods

Prospective observational study over 12 weeks of 252 patients with pain of any source lasting more than 6 months who sought integrative medicine evaluation and treatment at one of 9 Bravenet clinical sites across the United States. Interventions were not mandated. Baseline vitamin D status was reported to each clinician. Treatment options included conventional therapies as well as acupuncture, nutrition, massage and mind-body skills training. Measurements included the Brief Pain Inventory and 25-OH-vitamin D status.

Results

A total of 252 adults met eligibility criteria. Mean 25-OH-vitamin D levels and standard deviations at the start and 12 weeks later for all participants were 33.43 (17.05) and 39.58 (16.29). ($p < 0.0001$). The subset of low back pain patients demonstrated similar values of 32.71 (14.34) increasing after 12 weeks to 39.19 (13.26) ($p < 0.0001$). Of all participants, 153 (60.7%) achieved 25-OH-vitamin D levels above the 2010 international recommendation of ≥ 30 ng/ml and 99 (39.3%) did not. Of all 136 back pain participants, 90 (66.2%) achieved levels ≥ 30 ng/ml and 46 (33.8%) did not. Median average pain scores decreased from 5 to 4 during the study. Regardless of the integrative therapies applied, achievement of a serum level ≥ 30 ng/ml was necessary for significant improvement in average pain ($p = 0.0018$ for all patients, $p = 0.022$ for all back pain).

Conclusion

Chronic pain patients who sought integrative medicine demonstrated a high incidence of vitamin D deficiency. In the setting of additional treatments, serum levels ≥ 30 ng/ml were required for significant improvement in average pain scores. Failure to achieve a serum level ≥ 30 ng/ml represents a confounder of any therapeutic intervention in both clinical practice and clinical trials for chronic pain including chronic low back pain.

Published: 12 June 2012

doi:10.1186/1472-6882-12-S1-P120

Cite this article as: Plotnikoff et al.: P02.64. Vitamin D sufficiency is necessary for integrative treatment-associated improvements in chronic pain status. *BMC Complementary and Alternative Medicine* 2012 **12**(Suppl 1):P120.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



Penny George Institute for Health and Healing, Minneapolis, USA



© 2012 Plotnikoff et al; licensee BioMed Central Ltd. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.